


# Accepted Manuscript

## Genetic and Environmental Susceptibility to Food Allergy in a Registry of Twins

Juho E. Kivistö, MD, PhD, Ann Clarke, MD, PhD, Alizée Dery, Sarah De Schryver, MD, Greg Shand, MSc, Heini Huhtala, MSc, Mika J. Mäkelä, MD, PhD, Yuka Asai, MD, PhD, Kari Nadeau, MD, PhD, Laurie Harada, B.A., B.Ed, Edmond S. Chan, MD, Moshe Ben-Shoshan, MD, PhD



PII: S2213-2198(19)30471-4

DOI: <https://doi.org/10.1016/j.jaip.2019.05.016>

Reference: JAIP 2267

To appear in: *The Journal of Allergy and Clinical Immunology: In Practice*

Received Date: 1 April 2019

Revised Date: 3 May 2019

Accepted Date: 12 May 2019

Please cite this article as: Kivistö JE, Clarke A, Dery A, De Schryver S, Shand G, Huhtala H, Mäkelä MJ, Asai Y, Nadeau K, Harada L, Chan ES, Ben-Shoshan M, Genetic and Environmental Susceptibility to Food Allergy in a Registry of Twins, *The Journal of Allergy and Clinical Immunology: In Practice* (2019), doi: <https://doi.org/10.1016/j.jaip.2019.05.016>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Genetic and Environmental Susceptibility to Food Allergy in a Registry of Twins

Juho E. Kivistö<sup>1,2</sup>, MD, PhD; Ann Clarke<sup>3</sup>, MD, PhD; Alizée Dery<sup>4</sup>; Sarah De Schryver<sup>4</sup> MD; Greg Shand<sup>4</sup>, MSc.; Heini Huhtala<sup>5</sup>, MSc.; Mika J. Mäkelä<sup>6</sup>, MD, PhD; Yuka Asai<sup>7</sup> MD, PhD; Kari Nadeau<sup>8</sup> MD, PhD; Laurie Harada<sup>9</sup>, B.A., B.Ed; Edmond S. Chan<sup>10</sup>, MD; and Moshe Ben-Shoshan<sup>4</sup> MD, PhD

**Affiliations:** <sup>1</sup>Allergy Centre, Tampere University Hospital Tampere, Finland; <sup>2</sup>Pediatric Research Centre, University of Tampere, Finland; <sup>3</sup>Division of Rheumatology, Department of Medicine, Cumming School of Medicine, University of Calgary, Calgary, AB; <sup>4</sup>Division of Allergy and Clinical Immunology, Department of Pediatrics, McGill University, QC; <sup>5</sup>Faculty of Social Sciences, Tampere University, Finland; <sup>6</sup>Skin and Allergy Hospital, Helsinki University Hospital and University of Helsinki, Finland; <sup>7</sup>Division of Dermatology, Department of Medicine, Queen's University, Kingston, ON; <sup>8</sup>Department of Pediatrics, Division of Allergy, Immunology, and Rheumatology, Stanford University School of Medicine, CA; <sup>9</sup>Food Allergy Canada, ON; <sup>10</sup>Division of Allergy and Immunology, Department of Pediatrics, University of British Columbia, Vancouver, BC, Canada

**Address correspondence to:** Juho Kivistö, MD PhD, Allergy Centre, Tampere University Hospital, PL 2000, 33521 Tampere, Finland [juho.kivisto@tuni.fi], Tel: +358 40 530 4445, Fax: +358 3 215 6057

E-mail addresses for authors: [juho.kivisto@tuni.fi](mailto:juho.kivisto@tuni.fi), [aeclarke@ucalgary.ca](mailto:aeclarke@ucalgary.ca), [alizee.dery@gmail.com](mailto:alizee.dery@gmail.com), [sarah.deschryver@yahoo.com](mailto:sarah.deschryver@yahoo.com), [greg.shand@rimuhc.ca](mailto:greg.shand@rimuhc.ca), [heini.huhtala@tuni.fi](mailto:heini.huhtala@tuni.fi), [mika.makela@hus.fi](mailto:mika.makela@hus.fi), [yuka.asai@queensu.ca](mailto:yuka.asai@queensu.ca), [knadeau@stanford.edu](mailto:knadeau@stanford.edu), [lharada2018@gmail.com](mailto:lharada2018@gmail.com), [echan5@cw.bc.ca](mailto:echan5@cw.bc.ca), [moshebenshoshan@gmail.com](mailto:moshebenshoshan@gmail.com)

Study has been funded by the Tampere Tuberculosis Foundation, the Research Foundation of the Pulmonary Diseases, the Emil Aaltonen Foundation, Foundation of Tampere University Hospital, Foundation for Paediatric Research, Väinö and Laina Kivi Foundation, Finnish Medical Foundation, Allergy Research Foundation and The Finnish Society of Allergology and Immunology.

The authors do not have any additional affiliations, financial agreements, other involvement with any company, or any potential conflict of interest.

**Word count:** 1000

41

42

43 **Clinical Implications statement**

44 In a study of 80 twin pairs we demonstrate that genetic factors play a major role in the  
45 development of food allergy and that atopic dermatitis is a significant risk factor. Eczema  
46 control might reduce the risk of food allergy.

47

**To the Editor:**

A food allergy poses a substantial burden in many countries and the prevalence of food induced anaphylaxis is increasing<sup>1-3</sup>. It is likely that gene-environment interactions, rather than genetic factors solely, play a major role in the development of food allergy. Effective prevention of allergic diseases requires understanding of the factors that contribute to the development of allergy.

We aimed to evaluate the concordance rate for food allergy in pairs of MZ and DZ twins, for the most common food allergies. Moreover, we aimed to investigate the effect of zygosity, gender, co-morbidities and lifestyle habits on the development of food allergy.

Twins were recruited during 2014-2018 through Food Allergy Canada, Multiple Births Canada, BC Children's Hospital allergy clinic, and the Montreal Children's Hospital allergy clinic. Only participants with an allergist diagnosed food allergy AND the presence of convincing clinical history and positive confirmatory tests were included in this registry.

Interested participants were sent a consent form and a questionnaire, based on previous validated food allergy questionnaires<sup>4</sup> and the ISAAC (International Study of Asthma and Allergies in Childhood)<sup>5</sup>. The Principal Investigator and study coordinator independently reviewed participants' data.

DNA was collected through salivary samples, which were collected on all consenting and eligible participants to determine zygosity by genetic testing (GenePrint24 kit).

To assess twin concordance, we calculated probandwise concordance rates between pairs of MZ and DZ twins (defined as  $2C/(2C+D)$ ) where C is the number of all twin pairs that are both allergic to the specific food (concordant pairs), and D is the number of all discordant pairs). The probandwise rate is preferred over the pairwise rate as the probandwise concordance serves to forecast risk at the level of the individual rather than at the level of the pair. Further, pairwise concordance may underestimate the genetic effect<sup>6</sup>.

Univariable and multivariable logistic regression models were conducted to evaluate the association between genetic and environmental factors and the development of food allergy.

Statistical analysis was performed using R version 3.4.3 (2017-11-30). The McGill Research Ethics Boards approved the study (ethics reference number: 13-034 PED).

For this study, we recruited 80 twin pairs of which 34 were MZ, and 46 DZ. The median age of the patients was 4.8 years (range 0.59 – 35.8 years). Fifty-nine percent of the patients were boys and 41 % were girls.

Among 19 pairs of MZ and 30 pairs of DZ twins for peanut allergy, the concordance-rate was 0.59 and 0.29 respectively [difference= **0.31 (95%CI 0.04, 0.58)**]. Among 8 pairs of MZ and 8 pairs of DZ twins for pistachio allergy, the concordance-rate was 0.55 and 0.00 respectively [difference= **0.55 (95%CI 0.14, 0.95)**]. (Table 1)

Among 5 pairs of MZ and 6 pairs of DZ twins for walnut allergy, the concordance-rate was 0.57 and 0.00 respectively [difference= **0.57 (95%CI 0.05, 1.00)**]. Among 5 pairs of MZ and

4 pairs of DZ twins for sesame allergy the concordance-rate was 0.75 and 0.00 respectively [difference= **0.75 (95%CI 0.26, 1.00)**]. (Table 1)

When investigating the risk of allergy to any food, the odds ratio of the atopic dermatitis was 6.74 (**95%CI 2.29, 19.83, p=0.001**) in the univariable regression model and 6.41 (**95%CI 1.93, 21.28, p=0.02**) in the multivariable regression model when adjusted for gender, zygosity, atopic dermatitis and use of more than 4 courses of antibiotics. The same was observed for peanut allergy: odds ratio of the atopic dermatitis was 8.42 (**95%CI 2.09, 33.99, p=0.003**) in the univariable regression model and 8.3 (**95%CI 1.80, 38.27, p=0.007**) in multivariable regression model. (Table 2)

There was only one previous study on clinical food allergy (i.e. food allergy that was established through corroborating clinical symptoms of reaction with a positive confirmatory test) in twins. This study has shown higher concordance rate for peanut allergy among MZ twins compared to DZ twins (0.64 vs. 0.07)<sup>7</sup>. The present study shows similarly significant higher concordance rate of peanut allergy among MZ twins strengthening the evidence of heritability of peanut allergy. In addition, for the first time, we have shown a similar genetic effect among patients allergic to pistachio, walnut, sesame and fish. It is possible that genetic factors play more important role among certain tree nuts in the development of allergy.

Our study is unique as it identifies atopic dermatitis as a significant risk factor for food allergy, independent of genetic factors. This highlights the importance of atopic dermatitis control among children since this may reduce the risk of food allergy.

This study is novel, since this is the largest twin study evaluating the concordances of phenotyped food allergies among MZ and DZ twins. In one previous study, the sample size was larger but included only sensitization (not phenotyped food allergy)<sup>8</sup>. In another study, food allergy was based on parental report in contrast to our study when the presence of convincing history and positive confirmatory tests were required as well<sup>9</sup>. In addition, in contrast to previous studies, zygosity of the twin pairs was verified by genetic testing. The inclusion of all common food allergies is also a major strength of the study.

The present study has some limitations. First, some information may be subjected to recall bias. Second, some twins may have outgrown e.g. milk allergy. Moreover, the diagnosis was not confirmed by a food challenge in the majority of children. However, given that all cases were established by the presence of convincing history, confirmatory tests and allergist's diagnosis, we believe that any misclassification bias would be minimal. Finally, our sample size might have been too small to capture concordance differences and the effects of other factors on the risk of developing all major food allergies.

In summary, in this study including 80 twin pairs with median age of 5 years, we showed that genetic factors play a major role in the development of food allergies. This study showed that even when controlling for genetics, atopic dermatitis is a significant risk factor for food allergy. Further studies are needed to assess whether other risk factors (along with atopic dermatitis) will be identified as influencing the development of food allergies.

143 **Acknowledgements**

144 We would like to thank Multiple Birth Canada and Food Allergy Canada for helping  
145 with recruitment.

146

147



## References

1. Kivisto JE, Protudjer JL, Karjalainen J, Wickman M, Bergstrom A, Mattila VM. Hospitalizations due to allergic reactions in Finnish and Swedish children during 1999-2011. *Allergy* 2016; 71:677-83.
2. Lin RY, Anderson AS, Shah SN, Nurruzzaman F. Increasing anaphylaxis hospitalizations in the first 2 decades of life: New York State, 1990 -2006. *Ann Allergy Asthma Immunol* 2008; 101:387-93.
3. Protudjer JL, Jansson SA, Heibert Arnlind M, Bengtsson U, Kallstrom-Bengtsson I, Marklund B, et al. Household costs associated with objectively diagnosed allergy to staple foods in children and adolescents. *J Allergy Clin Immunol Pract* 2015; 3:68-75.
4. Ben-Shoshan M, Harrington DW, Soller L, Fragapane J, Joseph L, St Pierre Y, et al. A population-based study on peanut, tree nut, fish, shellfish, and sesame allergy prevalence in Canada. *J Allergy Clin Immunol* 2010; 125:1327-35.
5. Clayton T, Asher MI, Crane J, Ellwood P, Mackay R, Mitchell EA, et al. Time trends, ethnicity and risk factors for eczema in New Zealand children: ISAAC Phase Three. *Asia Pac Allergy* 2013; 3:161-78.
6. McGue M. When assessing twin concordance, use the probandwise not the pairwise rate. *Schizophr Bull* 1992; 18:171-6.
7. Sicherer SH, Furlong TJ, Maes HH, Desnick RJ, Sampson HA, Gelb BD. Genetics of peanut allergy: a twin study. *J Allergy Clin Immunol* 2000; 106:53-6.
8. Liu X, Zhang S, Tsai HJ, Hong X, Wang B, Fang Y, et al. Genetic and environmental contributions to allergen sensitization in a Chinese twin study. *Clin Exp Allergy* 2009; 39:991-8.
9. Ullemar V, Magnusson PK, Lundholm C, Zettergren A, Melen E, Lichtenstein P, et al. Heritability and confirmation of genetic association studies for childhood asthma in twins. *Allergy* 2016; 71:230-8.

Table 1. Specific food allergies concordances (C is the number of twin pairs that are both allergic to the specific food (concordant pairs), and D is the number of discordant pairs).

	Number of MZ pairs	Number of DZ pairs	Concordant pairs C	Discordant pairs D	MZ Concordance (2C/2C+D)	DZ Concordance (2C/2C+D)	Difference (95% Confidence Interval)
Almond	4	0	0	4	0.00		
Brazil nut	2	0	0	2	0.00		
Cashew	8	10	4	14	0.55	0.18	0.36 (-0.10, 0.82)
Codfish	1	1	1	1	1.00	0.00	<b>1.00 (0.25, 1.00)</b>
Egg	12	13	7	18	0.50	0.38	0.13 (-0.28, 0.53)
Fish	3	3	3	3	1.00	0.00	<b>1.00 (0.75, 1.00)</b>
Haddock	0	1	0	1		0.00	
Hazelnut	6	6	1	11	0.29	0.00	0.29 (-0.20, 0.78)
Kiwi	2	0	0	2	0.00		
Lentil	1	0	1	0	1.00		
Milk	5	4	1	8	0.33	0.00	0.33 (-0.25, 0.92)
Peanut	19	30	13	36	0.59	0.29	<b>0.31 (0.04, 0.58)</b>
Peas	1	0	1	0	1.00		
Pecan	5	2	1	6	0.33	0.00	0.33 (-0.38, 1.00)
Pinenut	0	1	0	1		0.00	
Pistachio	8	8	3	13	0.55	0.00	<b>0.55 (0.14, 0.09)</b>
Salmon	1	2	1	2	1.00	0.00	<b>1.00 (0.5, 1.00)</b>
Sesame	5	4	3	6	0.75	0.00	<b>0.75 (0.26, 1.00)</b>
Shellfish	4	1	1	4	0.40	0.00	0.40 (-0.43, 1.00)
Shrimp	2	0	2	0	1.00		
Soy	2	1	0	3	0.00	0.00	0.00
Sunflower	1	0	1	0	1.00		
Treenut	7	9	3	13	0.44	0.20	0.24 (-0.27, 0.76)
Trout	0	1	0	1		0.00	
Tuna	1	0	1	0	1.00		
Walnut	5	6	2	9	0.57	0.00	<b>0.57 (0.05, 1.00)</b>
Wheat	2	1	0	3	0.00	0.00	0.00
Pistachio/ Cashew	9	12	4	17	0.50	0.15	0.35 (-0.77, 0.07)
Walnut/ Pecan	6	6	2	10	0.50	0.00	<b>0.50 (0.01, 0.99)</b>
Any food	34	46	23	57	0.58	0.49	<b>0.10 (0.05, 0.46)</b>

Table 2. An association between genetic and environmental factors to the development of food allergy by logistic regression model.

Allergy			n of pairs	Univariate			Multivariable				
				OR	95% CI		P-value	OR	95% CI		p-value
Any food	Zygotity	di	46	1				1			
		mono	34	2.54	0.93	6.95	0.068	0.90	0.23	3.43	0.86
	Same gender	no	24	1				1			
		yes	56	3.60	0.95	13.59	0.059	3.17	0.56	18.05	0.19
	Eczema	(n)one	51	1				1			
		both	29	<b>6.74</b>	<b>2.29</b>	<b>19.83</b>	<b>0.001</b>	<b>6.41</b>	<b>1.93</b>	<b>21.28</b>	<b>0.02</b>
Peanut	Antibiotics	(n)one	76	1				1			
		both	4	9.00	0.88	91.76	0.064	9.99	0.88	113.80	0.064
	Zygotity	di	46	1				1			
		mono	34	2.52	0.74	8.55	0.137	0.65	0.14	3.15	0.60
	Same gender	no	24	1				1			
		yes	56	6.27	0.77	51.30	0.087	5.84	0.53	64.09	0.15
	Eczema	(n)one	51	1				1			
		both	29	<b>8.42</b>	<b>2.09</b>	<b>33.99</b>	<b>0.003</b>	<b>8.3</b>	<b>1.80</b>	<b>38.27</b>	<b>0.007</b>
	Introduction	(n)one	25	1				1			
		both	55	1.03	0.28	3.72	9.967	1.18	0.28	4.91	0.82
	Antibiotics	(n)one	76	1				1			
		both	4	1.31	0.14	12.79	0.815	0.93	0.079	10.77	0.95

Antibiotics = 4 or more courses of antibiotics, Introduction = Age of introduction to peanut (less than 1 year or more than 1 year)